

NYU Center for Data Science: DS-GA 1003  
Machine Learning and Computational Statistics (Spring 2019)

Brett Bernstein

February 27, 2019

**Instructions:** Following most lab and lecture sections, we will be providing concept checks for review. Each concept check will:

- List the lab/lecture learning objectives. You will be responsible for mastering these objectives, and demonstrating mastery through homework assignments, exams (midterm and final), and on the final course project.
- Include concept check questions. These questions are intended to reinforce the lab/lectures, and help you master the learning objectives.

You are strongly encourage to complete all concept check questions, and to discuss these (and related) problems on Piazza and at office hours. However, problems marked with a (★) are considered optional.

## Week 5 Lab: Concept Check Exercises

### Kernels

#### Kernel Learning Objectives

- Explain how explicit feature maps can be used to extend the expressivity of linear models.
- Explain potential issues explicitly computing large feature spaces.
- State and explain the definition of a 'kernelized' method.
- Explain why the SVM dual is kernelized, while the primal is not (ignoring the representer theorem).
- Give the relationship between a feature map and kernel function.
- Explain the computational benefits of kernelization based on costs of optimizing over  $\mathbb{R}^n$  vs  $\mathbb{R}^d$ .

- Be able to apply the kernel trick using the kernel matrix  $K$ .
- Be able to apply the elements of our proof of the representer theorem (ex: projections decrease norms) to prove related theorems.
- Compare using the representer theorem and duality to kernelized SVM.
- Describe common kernels (RBF/polynomial) and their properties (i.e. equivalent feature maps, computational benefits relative to explicit computation (if possible),...).
- Describe some general recipes for deriving "new" kernel function.

### Kernel Concept Check Questions

1. Fix  $n > 0$ . For  $x, y \in \{1, 2, \dots, n\}$  define  $k(x, y) = \min(x, y)$ . Give an explicit feature map  $\varphi : \{1, 2, \dots, n\}$  to  $\mathbb{R}^D$  (for some  $D$ ) such that  $k(x, y) = \varphi(x)^T \varphi(y)$ .
2. Show that  $k(x, y) = (x^T y)^4$  is a positive semidefinite kernel on  $\mathbb{R}^d \times \mathbb{R}^d$ .
3. Let  $A \in \mathbb{R}^{d \times d}$  be a positive semidefinite matrix. Prove that  $k(x, y) = x^T A y$  is a positive semidefinite kernel.
4. Consider the objective function

$$J(w) = \|Xw - y\|_1 + \lambda \|w\|_2^2.$$

Assume we have a positive semidefinite kernel  $k$ .

- (a) What is the kernelized version of this objective?
  - (b) Given a new test point  $x$ , find the predicted value.
5. Show that the standard 2-norm on  $\mathbb{R}^n$  satisfies the parallelogram law.
  6. Suppose you are given an training set of distinct points  $x_1, x_2, \dots, x_n \in \mathbb{R}^n$  and labels  $y_1, \dots, y_n \in \{-1, +1\}$ . Show that by properly selecting  $\sigma$  you can achieve perfect 0 – 1 loss on the training data using a linear decision function and the RBF kernel.
  7. Suppose you are performing standard ridge regression, which you have kernelized using the RBF kernel. Prove that any decision function  $f_\alpha(x)$  learned on a training set must satisfy  $f_\alpha(x) \rightarrow 0$  as  $\|x\|_2 \rightarrow \infty$ .
  8. Consider the standard (unregularized) linear regression problem where we minimize  $L(w) = \|Xw - y\|_2^2$  for some  $X \in \mathbb{R}^{n \times m}$  and  $y \in \mathbb{R}^n$ . Assume  $m > n$ .
    - (a) Let  $w^*$  be one minimizer of the loss function  $L$  above. Give an infinite set of minimizers of the loss function.
    - (b) What property defines the minimizer given by the representer theorem (in terms of  $X$ )?